

CORRECTION Open Access

Correction: The dynamics of vertebrate homeobox gene evolution: gain and loss of genes in mouse and human lineages

Ying-fu Zhong and Peter WH Holland*

Correction

After publication of this work [1], we found evidence for a gene (or two tandem genes) orthologous to *Leutx* in a genome assembly for the African elephant produced by the Broad Institute and accessible on the UCSC Genome Browser, assembly version Broad/loxAfr3 [2]. Elephant *Leutx* homeodomain sequences are highly

divergent from that of human *LEUTX* (48% identity), but regions of identity outside the homeodomain plus genomic synteny (Figure 1) confirm orthology. We also found divergent putative *Leutx* orthologues in some other mammals including dog, cat, horse, pig and cow but, as reported in our original publication, not in rodents or outside the placental mammals. These find-




Figure 1 The *LEUTX* homeobox gene (orange arrow) plus four unusual non-homeobox genes (light grey arrows) form a linked set of genes in the human genome (dashed box). Divergent putative *Leutx* genes are located in syntenic regions of cow, dog and elephant (orange arrows), but absent at the equivalent location in mouse and rat. Non-homeobox genes used as indicators of chromosomal synteny are shown as grey arrows.

^{*} Correspondence: peter.holland@zoo.ox.ac.uk Department of Zoology, University of Oxford, South Parks Road, Oxford, OX1 3PS, UK






Figure 2 A summary of homeobox gene dynamics in the mouse and human evolutionary lineages. The majority of homeobox genes are conserved between mouse and human lineages (grey squares), although some have undergone duplication to different extents (cascaded boxes). Humans have lost the *Msx3* gene; mice have lost *Leutx, Nanognb, Tprx1, LOC647589, Rax2, Shox, Dprx, Argfx* and *Ventx* (dashed boxes). Three new homeobox loci (*Gm7235, Gm5585* and *Crxos1*) and one new cluster (Obox) arose in the rodent lineage.

ings imply that *Leutx* did not originate in primate evolution as we originally suggested [1], but arose early in the radiation of placental mammals and was secondarily lost from rodents. This finding strengthens our original conclusion that rodents have experienced far more homeobox gene loss than have primates. *Leutx* is added to *Ventx, Argfx, Dprx, Shox, Rax2, LOC647589, Tprx1* and *Nanognb* on the list of homeobox genes lost in the evolutionary lineage leading to rodents (Figure 2).

Received: 8 July 2011 Accepted: 13 July 2011 Published: 13 July 2011

References

- Zhong Y-F, Holland PWH: The dynamics of vertebrate homeobox gene evolution: gain and loss of genes in mouse and human lineages. BMC Evolutionary Biology 2011, 11:169.
- 2. UCSC Genome Browser. [http://UCSC Genome Browser Home].

doi:10.1186/1471-2148-11-204

Cite this article as: Zhong and Holland: Correction: The dynamics of vertebrate homeobox gene evolution: gain and loss of genes in mouse and human lineages. *BMC Evolutionary Biology* 2011 11:204.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

